



Research note

Selected body temperature of nine species of Mexican horned lizards (*Phrynosoma*)

Temperatura corporal seleccionada de nueve especies mexicanas de lagartijas cornudas (Phrynosoma)

Rafael A. Lara-Reséndiz,^{a,b,*} Diego M. Arenas-Moreno,^{a,c}
Elizabeth Beltrán-Sánchez,^d Weendii Gramajo,^d Javier Verdugo-Molina,^e
Wade C. Sherbrooke,^f and Fausto R. Méndez-De la Cruz^a

^a Laboratorio de Herpetología, Departamento de Zoología, Instituto de Biología, Universidad Nacional Autónoma de México, Apartado postal 70515, 04510 Coyoacán, México, D. F., Mexico

^b Department of Ecology and Evolutionary Biology, Earth and Marine Sciences Building A316, University of California, Santa Cruz, 95064 California, USA

^c Posgrado en Ciencias Biológicas, Instituto de Biología, Universidad Nacional Autónoma de México, Apartado postal 70515, 04510 Coyoacán, México, D. F., Mexico

^d Instituto de Investigaciones Científicas, Área de Ciencias Naturales, Interior del Jardín Botánico, Universidad Autónoma de Guerrero, 39075 Chilpancingo, Guerrero, Mexico

^e Universidad de Sonora. Blvd. Luis Encinas S/N, Col. Centro, 83000 Hermosillo, Sonora, Mexico

^f Southwestern Research Station, American Museum of Natural History, P.O. Box 16553, Portal 85632, Arizona, USA

Received 17 June 2014; accepted 17 September 2014

Abstract

Temperature plays an important role in the behavior, ecology, physiology, and evolution of reptiles. Herein, we present selected body temperatures on 9 species of *Phrynosoma* under laboratory conditions, and note congeneric differences related to altitude and reproductive mode. The results are discussed with those of other phrynosomatid lizards.

All Rights Reserved © 2015 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access item distributed under the Creative Commons CC License BY-NC-ND 4.0.

Keywords: Thermal gradient; Phrynosomatids; Laboratory conditions; Preferred temperature; Reproductive mode; Altitude

Resumen

La temperatura juega un rol importante en el comportamiento, ecología, fisiología y evolución de reptiles. Aquí, se presentan las temperaturas corporales seleccionadas de nueve especies de *Phrynosoma* bajo condiciones de laboratorio, tomando en cuenta la relación con la altitud y el modo reproductor. Se discuten los resultados con las temperaturas corporales de otros phrynosomátidos.

Derechos Reservados © 2015 Universidad Nacional Autónoma de México, Instituto de Biología. Este es un artículo de acceso abierto distribuido bajo los términos de la Licencia Creative Commons CC BY-NC-ND 4.0.

Palabras clave: Gradiente térmico; Phrynosomátidos; Condiciones de laboratorio; Temperatura preferida; Modo reproductor; Altitud

* Corresponding author.

E-mail address: rafas.lara@gmail.com (R.A. Lara-Reséndiz).

In ectotherms most physiological mechanisms are affected by temperature, therefore they carefully regulate body temperature (T_b) within a relatively narrow range by using behavioral adjustments (Cowles & Bogert, 1944). T_b data collected in the field form the basis of most thermal biology reports in the literature (Avery, 1982). In contrast, selected body temperatures (T_{sel}) in laboratory conditions are rarely reported. T_{sel} , also known as preferred body temperature, represents the range of body temperatures at which an ectothermic animal seeks to maintain itself by behavioral means in order to carry on their normal activities (Brattstrom, 1965a). This temperature is commonly measured in laboratory conditions that provide an array of thermal environments free of physical and biotic constraints (e. g., predation, costs of thermoregulation; Clusella-Trullas, Terblanche, van Wyk, & Spotila, 2007). T_{sel} had been recorded for a wide range of diurnal lizard families, mostly in Scincidae, Agamidae, Lacertidae, Liolaemidae, and Phrynosomatidae (Clusella-Trullas & Chown, 2014), nonetheless, within this last family the T_{sel} of horned lizards (genus *Phrynosoma*) is actually quite rare in the literature (Lara-Reséndiz, Jezkova, Rosen, & Méndez-de la Cruz, 2014).

Horned lizards are one of the most distinctive lizards in North America. Most horned lizards have diets consisting primarily of native ants, dorsoventrally compressed bodies, cranial horns and body fringes, camouflaging coloration and sedentary habits, a constellation of unusual and interrelated traits including “relaxed” thermoregulation (Pianka & Parker, 1975). According to the latest taxonomic revision, this genus comprises 17 species distributed from Canada to Guatemala (Nieto-Montes de Oca, Arenas-Moreno, Beltrán-Sánchez, & Leaché, 2014). Among lizards, this genus has one of the largest altitudinal distributions (–200 to 3,440 m elev.) and 2 groups have evolved viviparity, apparently a characteristic associated with montane habitats (Hodges, 2004). There have been several reports on T_b in the field and critical temperatures in *Phrynosoma* (Cowles & Bogert, 1944; Heath, 1962; Lemos-Espinal, Smith, & Ballinger, 1997; Pianka & Parker, 1975;

Woolrich-Piña, Smith, & Lemos-Espinal, 2012, and others) and various other reports detailing aspects of its thermal ecology (Brattstrom, 1965b; Christian, 1998; Guyer & Linder, 1985; Powell & Russell, 1985; Prieto & Whitford, 1971, and others). Nevertheless T_{sel} of horned lizards has attracted relatively little study even though it is essential to understanding behavior, natural history, ecophysiology (Huey, 1982; Sherbrooke, 1997; Sherbrooke & Frost, 1989), and responses to the effects of global warming (Sinervo et al., 2010). Therefore, our principal objective was to determine the selected body temperatures of *Phrynosoma* in a laboratory thermal gradient and to discuss the results with those of other phrynosomatid lizards.

We conducted fieldwork during 4 years (2010-2013) in 9 localities from Mexico, A) Zumpago de Neri, Guerrero (*P. asio* and *P. taurus*; 17°36'11.73" N, 99°31'19.74" W, 1,130-1,400 m elev.) in a deciduous tropical forest; B) Ensenada, Baja California (*P. blainvilli*; 31°55'14.94" N, 116°36'32.6" W, 235 m elev.) in a chaparral community; C) volcán La Malinche, Tlaxcala (*P. orbiculare*; 19°14'38.49" N, 97°59'26.48" W, 3,140 m elev.) in a montane woodland and grassland; D) Chilapa de Álvarez, Guerrero (*P. sherbrookei*; 17°33'10.85" N, 99°16'15.01" W, 2,020 m elev.) in an oak forest with patches of grassland and scrub; E) Zapotitlán de las Salinas, Puebla (*P. taurus*; 18°19'39.68" N, 97°27'16.17" W, 1,400 m elev.) in a desert scrub with cacti and xerophilous plants; F) Hermosillo, Sonora (*P. solare*; 28°57'49.82" N, 111°4'1.2" W, 160 m elev.) in a desert scrub habitat, some with large, columnar cacti and small trees; G) Altamirano, Janos, Chihuahua (*P. hernandesii*; 30°20'58.24" N, 108°29'26.3" W, 1,830 m elev.) in an oak forest with patches of grassland and scrub; H) Samalayuca, Ciudad Juárez, Chihuahua (*P. modestum*; 31°28'33.1" N, 106°28'33.6" W, 1,260 m elev.) in sand dunes and semi stabilized dunes; I) Janos, Chihuahua (*P. cornutum*; 30°49'31.6" N, 108°28'26.6" W, 1,435 m elev.) in grasslands and scrublands.

All animals were captured by hand. Snout-vent length (SVL) was measured to the nearest 0.1 mm, and sex was determined

Table 1
Selected body temperature (T_{sel}) of horned lizards (*Phrynosoma*) and T_{sel} range (quartiles 25 and 75%) in °C. Localities description in text. SVL= snout-vent length, ND= not determined. *= Viviparous, **= oviparous

Species	n (♀:♂)	Adult body size (SVL; mm)	T_{sel}	T_{sel} 25%	T_{sel} 75%	Elevation (m asl)	References
<i>P. asio</i> **	67 (33:34)	≥90	35.9±0.13	34.3	38.0	1,260	Hodges, 2002; present study
<i>P. blainvilli</i> **	5 (3:2)	≥62	37.6±0.39	35.9	39.2	235	Goldberg, 1983; present study
<i>P. braconieri</i> *	78 (56:22)	≥49	35.9±ND	33.9	37.4	2,470	Zamudio and Parra-Olea, 2000; Hernández-Ríos, not published data
<i>P. cornutum</i> **	10	ND	38.5±ND	37.5	39.0	1,230	Prieto and Whitford, 1971
<i>P. cornutum</i> **	97 (61:36)	≥69	34.2±0.1	32.5	36.0	1,435	Lemos-Espinal and Smith, 2009; present study
<i>P. goodei</i> **	40 (16:24)	≥66	34.9±0.18	32.5	37.3	60	Lara-Reséndiz et al., 2014
<i>P. hernandesii</i> * (Chihuahua)	7 (6:1)	≥50	33.0±0.18	30.9	35.5	1,830	James, 2002; present study
<i>P. hernandesii</i> * (= <i>dougllassii</i>)	10	ND	35.0±ND	33.0	36.0	2,620	Prieto and Whitford, 1971
<i>P. modestum</i> **	8 (6:2)	≥37	33.6±0.51	31.1	36.5	1,260	Goldberg, 2012; present study
<i>P. orbiculare</i> * (Tlaxcala)	10 (5:5)	≥67	31.4±0.22	30.8	33.7	3,140	Hodges, 2002; present study
<i>P. orbiculare</i> * (State of México)	57	≥67	33.1±0.19	30.5	33.3	2,600	Hodges, 2002; Arias-Balderas, 2011
<i>P. platyrhinos</i> **	1	ND	36.0±0.14	ND	ND	1,440	Pianka and Parker, 1975
<i>P. sherbrookei</i> *	45 (26:19)	≥54	33.4±0.14	31.1	36.2	2,020	Nieto-Montes de et al., 2014; present study
<i>P. solare</i> **	26 (15:11)	≥69	37.5±0.12	36.8	38.5	160	Howard, 1974; present study)
<i>P. taurus</i> * (Puebla)	11 (8:3)	≥55.5	35.9±0.24	34.9	37.5	1,400	Zamudio and Parra-Olea, 2000; present study
<i>P. taurus</i> * (Guerrero)	23 (12:11)	≥55.5	34.3±0.23	32.4	36.6	1,130	Zamudio and Parra-Olea, 2000; present study




Figure 1. Mean selected body temperature (T_{sel}) of horned lizards (*Phrynosoma*) as a function of elevation (A) and reproductive mode (B).

for all individuals. Our data are based only on adult *Phrynosoma* (Table 1). In the laboratory, the lizards were maintained at 20 °C in plastic containers. Laboratory experiments were conducted one day after capture using a thermal gradient located in a 20 °C controlled-temperature room. The thermal gradient consisted of a wood box 150 cm × 100 cm × 70 cm high divided into 10 lengthwise tracks separated by insulation barriers to prevent behavioral influence of adjacent lizards. The substrate was 2-3 cm of sandy soil. At one end, and at the center of the box, 3 lamps of 100 W hanging at 30 cm to generate a thermal gradient from 20 to 50 °C throughout the tracks. The T_{sel} of individuals in the thermal gradient was taken manually each hour, using the digital thermometer Fluke model 51-II, according to the activity period observed in the field and previous records for each species. The 25% and 75% quartiles of each species' T_{sel} range (T_{sel25} and T_{sel75}) were calculated to obtain the lower and upper limits. After laboratory experiments, all lizards were released at the site of capture.

The results in Table 1 show the T_{sel} average ± standard error (mean ± SE) and sample size (n) of each species studied and previous published data. The means of T_{sel} of the *Phrynosoma* species reported here are within the range of T_{sel} observed in other species of *Phrynosoma* (Table 1). Furthermore, Sinervo et al. (2010) reported an average of T_b of 35.2 ± 0.20 °C for the family Phrynosomatidae considering 210 populations and a T_b range of 26.8-41.5 °C, which is similar to other diurnal lizards (Andrews, 1998). Similarly, Clusella-Trullas and Chown (2014) reported an average of T_{sel} of 35.1 ± 2.2 °C considering 20 species of phrynosomatids. Moreover, *Phrynosoma cornutum* had the highest T_{sel} (38.5 °C) and *P. orbiculare* had the lowest (31.4 °C) (Fig. 1). T_{sel}

of oviparous species varied from 33.5 to 38.5 °C and viviparous species from 31.4 to 35.9 °C (Fig. 1).

Although the T_{sel} have a genetic basis, in many cases this T_{sel} of lizards can be altered due to selective pressures of ambient temperatures, as an adaptation to habitat. This means that the T_{sel} of individuals may be influenced by local environment temperatures (Huey & Bennett, 1987; Sinervo, 1990). Many studies have addressed questions regarding labile or static aspects of lizard's thermal biology (see Clusella-Trullas & Chown, 2014 and references therein). Recently, it has been suggested that T_b and T_{sel} tend to be similar and conserved within closely related species, even at the level of genus, despite obvious morphological differences, such as size, coloration, reproductive mode or different environmental conditions (Grigg & Buckley, 2013). However, we found that the mean T_{sel} of horned lizard species seems to be related to the type of habitat, therefore not being overall conserved within the genus *Phrynosoma*. At localities with higher ambient temperatures horned lizard species tend to have higher average T_{sel} , while in cooler habitats horned lizard species have lower average T_{sel} . Again within the genus, higher and lower mean T_{sel} appear related, respectively, with oviparous and viviparous modes of reproduction (Fig. 1), as noted by Hodges (2004).

Future phylogenetic comparative analysis of thermal preferences of horned lizards should refine issues related to elevation, reproductive mode and other life history traits. In addition, the direction and magnitude of labile shifts in T_{sel} should be incorporated into mechanistic models (Lara-Reséndiz et al., 2014; Sinervo et al., 2010), to improve predictions of the impact of global climate change on populations of *Phrynosoma*.

In conclusion, the results presented above suggest that *Phrynosoma* maintain similar T_{sel} to other phrynosomatids and relatively higher T_{sel} at low elevations than at high elevations. However, this conclusion should be considered preliminary, pending broader data and incorporating biophysical models and phylogenetic analysis on their thermal biology.

This research was supported by a DGAPA-PAPIIT grant to F. R. Méndez-De la Cruz (IN215011-3). R. A. Lara-Reséndiz was funded by a Conacyt doctoral scholarship (165066) and a UC MEXUS-Conacyt postdoctoral fellowship (FE-14-2) to complete this manuscript. Collection permits (03699/09 and 05570/10) were given by Semarnat. We thank H. Gadsden, G. Castañeda, R. Huitrón, V. Jiménez, A. Díaz de la Vega, L. Rivera, S. González-Guzmán, L. López-Damián, M. García, D. Brindis, R. Santos, L. Florentino, and Solis-Ramírez family for field and laboratory assistance. Finally, we thank La Malinche Scientific Station, Helia Bravo Hollis Botanical Garden, and El Uno Ecological Reserve (TNC) for logistics support.

References

- Andrews, R. M. (1998). Geographic variation in field body temperature of *Sceloporus* lizards. *Journal of Thermal Biology*, 23, 329–334.
- Arias-Balderas, S. F. (2011). *Termorregulación y comportamiento de forrajero de Phrynosoma orbiculare en El Arenal, Nicolás Romero, Estado de México*. (M.Sc. Thesis). México, D.F.: Instituto de Biología, Universidad Nacional Autónoma de México.
- Avery, R. A. (1982). Field studies of body temperatures and thermoregulation. In C. Gans & F. H. Pough (Eds.), *Biology of the Reptilia* (pp. 93–166). New York: Academic Press.
- Brattstrom, B. H. (1965a). Body temperatures of reptiles. *American Midland Naturalist*, 73, 376–422.
- Brattstrom, B. H. (1965b). *Temperature regulation and diurnal activity in horned lizards*. Berkeley: University of California Publications in Zoology (pp. 97–136).
- Christian, K. A. (1998). Thermoregulation by the short-horned lizard (*Phrynosoma douglassi*) at high elevation. *Journal of Thermal Biology*, 23, 395–399.
- Clusella-Trullas, S., & Chown, S. L. (2014). Lizard thermal trait variation at multiple scales: a review. *Journal of Comparative Physiology B-Biochemical Systemic and Environmental Physiology*, 184, 5–21.
- Clusella-Trullas, S., Terblanche, J. S., van Wyk, J. H., & Spotila, J. R. (2007). Low repeatability of preferred body temperature in 4 species of cordylid lizards: temporal variation and implications for adaptive significance. *Evolutionary Ecology*, 21, 63–79.
- Cowles, R. B., & Bogert, C. M. (1944). A preliminary study of the thermal requirements of desert reptiles. *Bulletin of the American Museum of Natural History*, 83, 263–296.
- Goldberg, S. R. (1983). Reproduction of the coast horned lizard, *Phrynosoma coronatum*, in Southern California. *Southwestern Naturalist*, 28, 478–479.
- Goldberg, S. R. (2012). Notes on reproduction in roundtail horned lizards, *Phrynosoma modestum* (Squamata: Phrynosomatidae) from New Mexico. *Bulletin of the Chicago Herpetological Society*, 47, 34–36.
- Grigg, J. W., & Buckley, L. B. (2013). Conservatism of lizard thermal tolerances and body temperatures across evolutionary history and geography. *Biology Letters*, 9, 20121056.
- Guyer, C., & Linder, A. D. (1985). Thermal ecology and activity patterns of the short-horned lizard (*Phrynosoma douglassi*) and the sagebrush lizard (*Sceloporus graciosus*) in southeastern Idaho. *Great Basin Naturalist*, 45, 607–614.
- Heath, J. E. (1962). Temperature-independent morning emergence in lizards of the genus *Phrynosoma*. *Science*, 138, 891–892.
- Hodges, W. L. (2002). *Phrynosoma systematics, comparative reproductive ecology, and conservation of a Texas native*. (Ph.D. Thesis). Austin: University of Texas.
- Hodges, W. L. (2004). Evolution of viviparity in horned lizards (*Phrynosoma*): testing the cold-climate hypothesis. *Journal of Evolutionary Biology*, 17, 1230–1237.
- Howard, C. W. (1974). Comparative reproductive ecology of horned lizards (Genus *Phrynosoma*) in southwestern United States and northern Mexico. *Journal of the Arizona Academy of Science*, 9, 108–116.
- Huey, R. B. (1982). Temperature, physiology, and ecology of reptiles. In C. Gans & F. H. Pough (Eds.), *Biology of the Reptilia* (pp. 25–91). New York: Academic Press.
- Huey, R. B., & Bennett, A. F. (1987). Phylogenetic studies of coadaptation: preferred temperatures versus optimal performance temperatures of lizards. *Evolution*, 41, 1098–1115.
- James, J. D. (2002). A survey of short-horned lizard (*Phrynosoma hernandesi hernandesi*) populations in Alberta. *Alberta Species at Risk Report*, 29, 1–25.
- Lara-Reséndiz, R. A., Jezkova, T., Rosen, P. C., & Méndez-de la Cruz, F. R. (2014). Thermoregulation during the summer season in the Goode's horned lizard *Phrynosoma goodie* (Iguania: Phrynosomatidae) in Sonoran Desert. *Amphibia-Reptilia*, 35, 161–172.
- Lemos-Espinal, J. A., & Smith, H. M. (2009). *Anfibios y reptiles del estado de Chihuahua, México/Amphibians and reptiles of the state of Chihuahua, Mexico*. México, D. F.: Universidad Nacional Autónoma de México-Comisión para el Conocimiento y Uso de la Biodiversidad.
- Lemos-Espinal, J. A., Smith, G. R., & Ballinger, R. E. (1997). Observations on the body temperatures and natural history of some Mexican reptiles. *Bulletin of the Maryland Herpetological Society*, 33, 159–164.
- Nieto-Montes de Oca, A., Arenas-Moreno, D., Beltrán-Sánchez E., & Leaché, A. (2014). A new species of horned lizard (genus *Phrynosoma*) from Guerrero, México, with an updated multilocus phylogeny. *Herpetologica*, 70, 241–257.
- Pianka, E. R., & Parker, W. S. (1975). Ecology of horned lizards: review with special reference to *Phrynosoma platyrhinos*. *Copeia*, 1975, 141–162.
- Powell, G. L., & Russell, A. P. (1985). Field thermal ecology of the eastern short-horned lizard (*Phrynosoma douglassi brevirostre*) in southern Alberta. *Canadian Journal of Zoology*, 63, 228–238.
- Prieto, A. A., Jr., & Whitford, W. G. (1971). Physiological responses to temperature in horned lizards, *Phrynosoma cornutum* and *Phrynosoma douglassii*. *Copeia*, 1971, 498–504.
- Sherbrooke, W. C. (1997). Physiological (rapid) change of color in horned lizards (*Phrynosoma*) of arid habits: hormonal regulation, effects of temperature, and role in nature. *Amphibia-Reptilia*, 18, 155–175.
- Sherbrooke, W. C., & Frost, S. K. (1989). Integumental chromatophores of a color-change thermoregulating lizard *Phrynosoma modestum* (Iguanidae; Reptilia). *American Museum Novitates*, 2943, 1–14.
- Sinervo, B. (1990). Evolution of thermal physiology and growth rate between populations of the western fence lizard (*Sceloporus occidentalis*). *Oecologia*, 83, 228–237.
- Sinervo, B., Méndez-de la Cruz, F., Miles, D. B., Heulin, B., Bastiaans, E., Villagrán-Santa Cruz, M., et al. (2010). Erosion of lizard diversity by climate change and altered thermal niches. *Science*, 328, 894–899.
- Woolrich-Piña, G. A., Smith, G. R., & Lemos-Espinal, J. A. (2012). Body temperatures of 3 species of *Phrynosoma* from Puebla, Mexico. *Herpetology Notes*, 5, 361–364.
- Zamudio, K. R., & Parra-Olea, G. (2000). Reproductive mode and female reproductive cycles of 2 endemic Mexican horned lizards (*Phrynosoma taurus* and *Phrynosoma braconieri*). *Copeia*, 2000, 222–229.