



Available online at www.sciencedirect.com

Revista Mexicana de Biodiversidad

Revista Mexicana de Biodiversidad 87 (2016) 1149–1152



www.ib.unam.mx/revista/

Research note

New occurrence records of *Eleutherodactylus syristes* in Guerrero and Oaxaca, Mexico

Nuevos registros de distribución de *Eleutherodactylus syristes* en Guerrero y Oaxaca, México

Uri Omar García-Vázquez^{a,*}, Thomas J. Devitt^{b,1}, Adrián Nieto-Montes de Oca^c, Sean M. Rovito^{b,2}, Itzue Caviedes-Solis^c, Gabriela Parra-Olea^b

^a Facultad de Estudios Superiores Zaragoza, Universidad Nacional Autónoma de México, Batalla 5 de Mayo, Ejército de Oriente, 09230 Ciudad de México, Mexico

^b Departamento de Zoología, Instituto de Biología, Universidad Nacional Autónoma de México, Apartado postal 70-153, 04510 Ciudad de México, Mexico

^c Laboratorio de Herpetología y Museo de Zoología, Facultad de Ciencias, Universidad Nacional Autónoma de México, Circuito Exterior s/n, 04510 Ciudad de México, Mexico

Received 3 November 2015; accepted 4 April 2016

Available online 7 August 2016

Abstract

The Sierra Madre del Sur (SMS) of southern Mexico is known for its exceptionally high levels of amphibian endemism, but knowledge of species' ranges in the region is incomplete. Here, we report new locality records for *Eleutherodactylus syristes* (Hoyt, 1965) from the states of Oaxaca and Guerrero. Previously, this species was known only from the Pacific slopes of the Sierra de Miahuatlán and Mixteca Alta in Oaxaca. These new occurrence records extend the known range of this species by approximately 335 km from the type locality and increase the elevational range, indicating it is more widespread in the SMS. *Eleutherodactylus syristes* is currently listed as endangered by the IUCN based on its limited distribution and ongoing threats to its persistence.

All Rights Reserved © 2016 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access item distributed under the Creative Commons CC License BY-NC-ND 4.0.

Keywords: Anura; Geographic distribution; Sierra Madre del Sur; *Syrrhopus*

Resumen

La sierra Madre del Sur en México es reconocida por su alto número de anfibios endémicos. Sin embargo, el conocimiento de la distribución de estas especies es incompleto. En este trabajo registramos localidades nuevas de *Eleutherodactylus syristes* (Hoyt, 1965) para los estados de Oaxaca y Guerrero. Esta especie era conocida solo para la vertiente del Pacífico de la sierra de Miahuatlán y para la Mixteca alta en Oaxaca. Los nuevos registros extienden la distribución conocida de esta especie a aproximadamente 335 km de la localidad tipo y aumentan su intervalo altitudinal, indicando que su distribución en la sierra Madre del Sur es más extendida. *Eleutherodactylus syristes* está considerada actualmente en peligro por la UICN con base en su limitada distribución geográfica y las amenazas actuales para su persistencia.

Derechos Reservados © 2016 Universidad Nacional Autónoma de México, Instituto de Biología. Este es un artículo de acceso abierto distribuido bajo los términos de la Licencia Creative Commons CC BY-NC-ND 4.0.

Palabras clave: Anura; Distribución geográfica; Sierra Madre del Sur; *Syrrhopus*

* Corresponding author.

E-mail address: urigarcia@gmail.com (U.O. García-Vázquez).

¹ Current address: Department of Integrative Biology, The University of Texas at Austin, Austin, TX 78712, USA.

² Current address: LANGEBIO-CINVESTAV, Km 9.6 Libramiento Norte Carretera León, 36821 Irapuato, Guanajuato, Mexico.

Peer Review under the responsibility of Universidad Nacional Autónoma de México.

Eleutherodactyline frogs of the subgenus *Syrrhophus* are some of the most ubiquitous and abundant of Mexican amphibians, yet also among the least studied. These terrestrial, direct-developers occur at low to moderate elevations (up to 2500 m) from Texas through Mexico and into Guatemala and Belize, with 2 species in western Cuba (Hedges, Duellman, & Heinicke, 2008). Several species have been documented only from the type locality (IUCN, 2016). *Eleutherodactylus syristes* was described by Hoyt (1965) from 62.5 to 66.7 km (by road) N of Pochutla, Oaxaca, Mexico between 850 and 1620 m in elevation. Santos-Barrera and Canseco-Márquez (2004) reported this species from the Mixteca Alta in Oaxaca, but did not provide any additional information.

As part of ongoing revisionary work on this group, we conducted fieldwork throughout Mexico from 2007 to 2013, including several localities from Guerrero and Oaxaca in the Sierra Madre del Sur (SMS). This region is known for its exceptionally high levels of amphibian endemism (Campbell, 1999; Delia, Whitney, & Burkhardt, 2013; Flores-Villela, Canseco-Márquez, & Ochoa-Ochoa, 2010). A well-known locality that appears often in the literature on amphibians and reptiles inhabiting the SMS is Agua del Obispo, Guerrero (Campbell & Duellman, 2000; Davis & Dixon, 1955, 1964; Dixon, 1957; Lips, Mendelson, Muñoz-Alonso, Canseco-Márquez, & Mulcahy, 2004; Mendelson, Williams, Sheil, & Mulcahy, 2005; Taylor, 1940). Agua del Obispo represents the type locality for several

species, including *Eleutherodactylus albolabris* (Taylor, 1943). Located on the old Mexico City-Acapulco highway (MX95) approximately 31 km south of Chilpancingo, Agua del Obispo does not appear on maps because it is a small, private hacienda (Mendelson et al., 2005). Elevation is approximately 1000 m. Habitat there consists of scattered small pines and brush in an ecotone between tropical deciduous forest at lower elevations and pine-oak woodland at higher elevations (Campbell & Duellman, 2000).

On 9 September 2010, we visited Agua del Obispo in search of *E. albolabris*. A brief search yielded a single specimen of *Eleutherodactylus*, which could not be assigned to *E. albolabris*. The specimen was collected under some roofing tiles on the forest floor, and has been deposited in the Museo de Zoología, Facultad de Ciencias, Universidad Nacional Autónoma de México (MZFC 28683). This specimen fits the description of another specimen from Agua del Obispo deposited in the Texas Cooperative Wildlife Collection at Texas A&M University (TCWC 11261) collected by Ralph W. Axtell and reported by Davis and Dixon (1955). These authors tentatively assigned TCWC 11261 to *Tomodactylus nitidus petersi* (=*Eleutherodactylus nitidus*), but Dixon (1957) later assigned it to *T. albolabris* based on overall proportions and ventral color pattern. In his description of *T. syristes*, Hoyt (1965) discussed TCWC 11261, noting that this specimen resembled *T. syristes* more than *E. albolabris*. He suggested that it might be




Figure 1. Known distribution of *Eleutherodactylus syristes*.



Figure 2. Photographs of *Eleutherodactylus syristes*. Photos of MZFC 26863, MZFC 21686, and ANMO 1992, and remaining specimens taken by TJD, UOGV, and Peter Heimes, respectively.

conspecific with *T. syristes*, but did not assign it to any named species.

We have examined TCWC 11261 and find that both this specimen and MZFC 28683 are assignable to *E. syristes*, which had not been recorded previously from the state of Guerrero (Pérez-Ramos, Saldaña, & Uribe-Peña, 2000). These specimens extend the known range of this species approximately 335 km to the northwest from the type locality in the Sierra de Miahuatlán (Hoyt, 1965). In addition to Agua del Obispo, we have recorded this species at 6 other geographically intermediate localities in Guerrero and Oaxaca (Fig. 1; Appendix). These records extend the known elevational range of this species to 2320 m.

All specimens exhibit the diagnostic characters described by Hoyt (1965), including tips of the outer 2 fingers slightly expanded and rounded; dorsum smooth; in life, dark gray with indistinct black mottling; venter with extensive black mottling; and nearly all of anterior, dorsal, and posterior surfaces of thighs bright orange (Fig. 2). Additional morphological and color

pattern characters that distinguish this species from congeners are provided by Hoyt (1965).

Currently, 7 of 24 species of mainland *Eleutherodactylus* (*Syrrhophus*) are classified as endangered or critically endangered according to the IUCN Red List of Threatened Species criteria, including *E. syristes* (IUCN, 2016). Although the new distribution records presented here indicate that this species' range is more extensive in the SMS of Guerrero and Oaxaca, ongoing habitat loss and fragmentation throughout the SMS continue to threaten its persistence.

Fieldwork was supported by Conacyt grant number 154093 to A. Nieto-Montes de Oca and UC MEXUS-Conacyt postdoctoral fellowships to T. Devitt and S. Rovito. We thank the following persons for help with fieldwork: E. Centenero, P. Heimes, R. Meza, E. Pérez, and J.J. Avendaño. We also thank J.C. Arenas for assistance with the map. Collecting permits were granted by the Secretaría de Medio Ambiente y Recursos Naturales (Semarnat) to Gabriela Parra-Olea (FAUT-0106) and Uri Omar García-Vazquez (FAUT-0274).

Appendix. Specimens examined

- Eleutherodactylus syristes*.—México:
- Guerrero (8): Chilpancingo de los Bravo: Agua del Obispo (TCWC 11261, MZFC 28683). Malinaltepec: Road San Luis Acatlán to Malinaltepec, between San Luis Acatlán and Tres Marías (MZFC 28679-82); Paraje Montero (MZFC 28676-78).
- Oaxaca (21): Pluma Hidalgo: Finca Cruz Grande (MZFC 22506); Road to Jordán (MZFC 22555, 22609); Finca El Carmen (MZFC 22492); Río Copalita (MZFC 22525); Pluma Hidalgo (MZFC 22469, 22500-01). Putla Villa de Guerrero: Road between Putla and Pinotepa, 12.3 km S of Putla (MZFC 23938). San Augustín Loxicha: 24.5 km N of Candelaria Loxicha (MZFC 15856). San Andrés Cabecera Nueva: Guadalupe Zacatepec (MZFC 13390-91, 13395); Corral de Piedra, Santa Ana del Progreso (MZFC 13389, 13392-94). San Juan Lachao: Road between San Juan Lachao and Cerro de Vidrio, approx. 8.2 km S of turn-off to Santa Catarina Jiquila (MZFC 21686-7). Santa Lucía Monte Verde: 1 km N la Pedrera (ANMO 1992). Santo Domingo Yanhuitan: Road Huajapan-Nochixtlán Km 68 (MZFC 21655).
- References**
- Campbell, J. A. (1999). Distribution patterns of amphibians in Middle America. In W. E. Duellman (Ed.), *Patterns of distribution of amphibians* (pp. 111–210). Baltimore: The Johns Hopkins University Press.
- Campbell, J. A., & Duellman, W. E. (2000). New species of stream-breeding hylid frogs from the northern versant of the highlands of Oaxaca, Mexico. *Scientific Papers, Natural History Museum, The University of Kansas*, 16, 1–28.
- Davis, W. B., & Dixon, J. R. (1955). Notes on Mexican toads of the genus *Tomodactylus* with the descriptions of two new species. *Herpetologica*, 11, 154–160.
- Davis, W. B., & Dixon, J. R. (1964). Amphibians of the Chilpancingo Region, México. *Herpetologica*, 20, 225–233.
- Delia, J. R. J., Whitney, J. L., & Burkhardt, T. (2013). Rediscovery of 'lost' treefrogs from the Oaxacan highlands of Mexico. *Biodiversity and Conservation*, 22, 1405–1414.
- Dixon, J. R. (1957). Geographic variation and distribution of the genus *Tomodactylus* in Mexico. *Texas Journal of Science*, 9, 379–409.
- Flores-Villela, O., Canseco-Márquez, L., & Ochoa-Ochoa, L. M. (2010). Geographic distribution and conservation of the Mexican central highlands Herpetofauna. In L. D. Wilson, & J. H. Townsend (Eds.), *Conservation of mesoamerican amphibians and reptiles* (pp. 302–321). Eagle Mountain, Utah: Eagle Mountain Publishing, LC.
- Hedges, S. B., Duellman, W. E., & Heinicke, M. P. (2008). New World direct-developing frogs (Anura: Terrarana): molecular phylogeny, classification, biogeography, and conservation. *Zootaxa*, 1737, 1–182.
- Hoyt, D. L. (1965). A new frog of the genus *Tomodactylus* from Oaxaca. *Journal of the Ohio Herpetological Society*, 5, 19–22.
- IUCN (2016). The IUCN Red List of Threatened Species. Retrieved on March 6th 2016, from: <http://www.iucnredlist.org>.
- Lips, K. R., Mendelson, J. R., III, Muñoz-Alonso, A., Canseco-Márquez, L., & Mulcahy, D. G. (2004). Amphibian population declines in montane southern Mexico: resurveys of historical localities. *Biological Conservation*, 119, 555–564.
- Mendelson, J. R., Williams, B., Sheil, C., & Mulcahy, D. G. (2005). Systematics of the *Bufo coccifer* complex (Anura: Bufonidae) of Mesoamerica. *Scientific Papers, Natural History Museum, The University of Kansas*, 38, 1–27.
- Pérez-Ramos, E., Saldaña, L., & Uribe-Peña, Z. (2000). A checklist of the reptiles and amphibians of Guerrero, Mexico. *Anales del Instituto de Biología, Universidad Nacional Autónoma de México, Serie Zoología*, 71, 21–40.
- Santos-Barrera, G., & Canseco-Márquez, L. (2004). *Eleutherodactylus syristes*. The IUCN Red List of Threatened Species 2004: e.T56989A11552730. Retrieved on: October 18, 2015 from: <http://dx.doi.org/10.2305/IUCN.UK.2004.RLTS.T56989A11552730>.
- Taylor, E. H. (1940). Herpetological miscellany No. I. *University of Kansas Science Bulletin*, 26, 489–571.
- Taylor, E. H. (1943). Herpetological novelties from Mexico. *University of Kansas Science Bulletin*, 29, 343–358.