





Figures 1-3. Male holotype of *Isaeus neglectus* (BMNH). Dorsal, ventral and lateral view, respectively. Scale: 2.0 mm.



Figures 4-6. Males of *Isaeus neglectus* (CNAN). 4 and 5, dorsal habitus from Tenejapan and Sótano desbloqueado y Sótano de la fertilidad, respectively. 6, lateral habitus from Sótano de la fertilidad y Sótano desbloqueado. Scale: 2.0 mm.








Figures 7, 8. Chelicera of male of *Isaeus neglectus* (CNAN). Frontal and mesal view, respectively. Scale: 1.00 mm.



Figures 9-11. Pedipalp of male of *Isaeus neglectus* (CNAN). Ectal, mesal and frontal view, respectively. Scale: 0.9 mm.




Figures 12-14. Posterior legs of the male of *Isaeus neglectus* (CNAN). 12, mesal view of leg III.
13 and 14, ventral view of femora III and IV, respectively. Scale: 1.5 mm.



Figures 15-19. Male genitalia of *Isaeus neglectus* (CNAN). 15 and 16, frontal and lateral view of pars distalis. 17, detailed lateral view. 18 and 19, close-up of apical follis. Scale: 15, 16 = 150 μm , 17 = 50 μm , 18, 19 = 25 μm .



Figures 20, 21. Female holotype of *Isaeus phillippii* (SMF RII 2511/66). Dorsal and lateral view, respectively. Scale: 2.0 mm.



Figures 22, 23. Chelicera and pedipalp of the female holotype of *Isaeus phillippii* (SMF RII 2511/66). 22, frontal view of chelicerae. 23, mesal view of pedipalp. Scale: 1.0 mm.